

Publikationen von Dr. Alexandra H. Müller-Öffner

Socioeconomic studies of haemostaseologic therapy and anticoagulation

Schramm W, **Offner A**, Szucs T.

Zentrabl Chir. 1995; 120 (8): 593-7.

Socioeconomic impact of haemophilia care: results of a pilot study.

Szucs TD, **Öffner A**, Schramm W.

Haemophilia; 1996; 2 (4): 211-7.

Autoradiographic visualization of B1 bradykinin receptors in porcine vascular tissues in the presence or absence of inflammation.

Schremmer-Danninger E, **Offner A**, Siebeck M, Heinz-Erian P, Gais P, Roscher AA.

Immunopharmacology. 1996; 33 (1-3): 95-100.

Resource utilisation in haemophiliacs treated in Europe: results from the European Study on Socioeconomic Aspects of Hemophilia Care.

Szucs TD, **Offner A**, Kroner B, Giangrande P, Berntorp E, Schramm W.

Haemophilia 1998; 4 (4), 498-501.

B1 bradykinin receptors and carboxypeptidase M are both upregulated in the aorta of pigs after LPS-infusion.

Schremmer-Danninger E, **Offner A**, Siebeck M, Roscher AA.

Biochem Biophys Res Commun. 1998; 243;(1) 246-52.

Das metabolische Syndrom bei polyzystischen Ovarien: Ätiologie, Diagnose und Therapie im Überblick.

Janssen OE, **Öffner A**, Schopohl J, Brabant G.

Klinikarzt. 2000; 6: 163-6.

Quality of life, psychosocial well-being and sexual satisfaction in woman with polycystic ovary syndrome.

Elsenbruch S, Hahn S, Kowalsky D., **Offner AH**, Schedlowski M, Mann K, Janssen OE.

J Clin Endocrinol Metab. 2003; 88(12): 5801-7.

High prevalence of autoimmune thyroiditis in patients with polycystic ovary syndrome.

Janssen OE, Mehlmauer N, Hahn S., **Offner AH**, Gärtner R.

Eur J Endocrinol. 2004; 150 (3): 363-9.

Stenting of the superficial femoral artery after suboptimal ballon angioplasty: one-year results.

Abahji TN, Tato F, Rieger J., **Offner A**, Will S, Hoelscher4 G., Weiss N, Hoffmann U.

Int Angiol. 2006; 25 (2): 184-9.

Comparison of insulin glargine versus NPH insulin in people with Type 2 diabetes mellitus under outpatient-clinic conditions for 18 months using a basal-bolus regimen with a rapid-acting insulin analogue as mealtime insulin.

Siegmund T, Blankenfeld H, **Oeffner A**, Schumm-Draeger PM.
Exp Clin Endocrinol Diabetes. 2007; 115 (6): 349-53

Are symptoms of depression more common in diabetes? Results from the Heinz Nixdorf Recall Study.

Icks A, Krus J, Dragano N, Broecker-Preuss M, Slomiany U, Mann K, Jöckel KH, Erbel R, Giani G, Moebus S; **Heinz Nixdorf Recall Study Investigator Group**
Diabetes Med. 2008; 25 (11): 1330-6

Individuelle Therapie bei Diabetes mellitus: Diagnostik und Therapie des diabetischen Fußsyndroms.

Müller-Öffner A, Schumm-Draeger PM.
Notfall & Hausarztmedizin. 2007; 33 (3): 146-151.

Prevalence of anterior pituitary dysfunction in patients following traumatic brain injury in a German multi-centre screening program.

Berg C, **Oeffner A**, Schumm-Draeger PM, Badorrek F, Brabant G, Gerbert B, Bornstein S, Zimmermann A, Weber M, Broecker-Preuss M, Mann K, Herrmann BL. Exp Clin Endocrinol Diabetes. 2010; 118 (2): 139-44.

Coronary artery calcification and its relationship to validated genetic variants for diabetes mellitus assessed in the Heinz Nixdorf recall cohort.

Pechlivanis S, Scherag A, Mühleisen TW, Möhlenkamp S, Horsthemke B, Boes T, Bröcker-Preuss M, Mann K, Erbel R, Jöckel KH, Nöthen MM, Moebus S; **Heinz Nixdorf Recall Study Group**
Arterioscler Thromb Vasc Biol. 2010; 30 (9): 1867-72

Are air pollution and traffic noise independently associated with atherosclerosis: the Heinz Nixdorf Recall Study

Kälsch H, Hennig F, Moebus S, Möhlenkamp S, Dragano N, Jakobs H, Memmesheimer M, Erbel R, Jöckel KH, Hoffmann B; **Heinz Nixdorf Recall Study Investigative Group**.
Eur Heart J. 2014 Apr;35(13):853-60.